

THE PPDB

Background and Support Information

The present document aims at providing information regarding the PPDB, its management and data acquisition. It seeks to answer common questions regarding these matters. Information on the Terms and Conditions of use can be found in a document 'PPDB conditions of Use' that is available on the PPDB website as a PDF download.

Contents

	Page number
What is the Pesticide Properties Database?	1
How does the PPDB work?	1
Explanation of the Terms and Parameters used	2
1. General chemical status	2
2. Product and branding information	4
3. Properties related to environmental fate	4
4. Metabolites/Parents	7
5. Ecotoxicology	7
6. Human health	8
7. Translations	9
Data Sources used to Populate the PPDB	9
Data Interpretation	11
1. Environmental fate	11
2. Ecotoxicology	12
3. Human health	13
4. Notes	14
Key Species and Endpoint Glossary	17
Quality Control and Management	18

What is the Pesticide Properties Database?

The Pesticide Properties Database (PPDB) is a comprehensive relational database of pesticide physicochemical, toxicological, ecotoxicological and other related data.

The database has been developed by the Agriculture & Environment Research Unit (AERU) based at the University of Hertfordshire, UK. It has evolved from a database that originally accompanied the EMA (Environmental Management for Agriculture) software (also developed by AERU), with additional input from the EU-funded FOOTPRINT project (<http://www.eu-footprint.org>).

How does the PPDB work?

The data is collated and stored in an MS Access database. A Microsoft Visual Basic data management application has been developed to manage the data adhering to strict formatting and quality control protocols and to keep track of versions, literature references and record

cross-referencing. Another Visual Basic application is used to filter the data, handle the translations and generate the thousands of HTML pages for the PPDB website.

Explanation of the Terms and Parameters Used

(1) General chemical status information – descriptors and registration

Parameter	Explanation
Alias's/synonyms:	Other names by which the pesticide is known. These are enabled in the PPDB search facility.
Description:	General description of the major uses of the pesticide active substance.
Introduction:	Year (and country where known) of introduction, registration or discovery.
EU approval:	Provides an indication of where the active substance has been authorised for agricultural use within the EU. Please check with the relevant national authority before relying on this data.
EC Directive status:	Status of the chemical in the EU peer review process (91/414) of pesticide active substances. 1st stage - Data list, 2nd stage - Dossier, 3rd stage - Conclusion (Annex 1 or Exclusion or other status).
Also registered in:	Other countries where we believe the substance is used. Please check with the relevant national authority before relying on this data.
Pesticide type:	The specific type of pesticides described according to the type of pest they control e.g. Insecticide, Herbicide, Fungicide, Acaricide etc.
Chemical group:	Chemical classification group based on the chemical structure.
Mode of action:	The mechanism by which the pesticide performs its main function.
Metabolite type:	General description of the host process that creates the metabolite e.g. soil, water, animal, plant, groundwater.
Other constituent type:	General description of the purpose of the constituent in the formulation i.e. solvent, wetter, carrier etc.
CAS RN:	Chemical Abstracts Service Registry Number - a unique identify for the chemical.
EC number:	The unique reference number for the pesticide in the European Chemical Substances Information System (EINECS) or European List of Notified Chemicals (ELINCS).
CIPAC number:	The CIPAC code number system is a simple approach for an unambiguous coding of active ingredients and other substances used in the area/field of pesticides. CIPAC, FAO, WHO and the EU are the main users of this system.
US EPA chemical code:	The U.S. Environmental Protection Agency (U.S. EPA) assigns a unique reference number to individual pesticide active ingredients to assist in their identification. This code is sometimes referred to as the Shaugnessy Number.
Chemical formula:	This is a concise way of expressing information about the atoms that constitute the chemical.
SMILES:	The S implified M olecular I nput L ine E nter S pecification (SMILES) is a specification for describing the structure of chemical molecules using short ASCII strings. SMILES strings can be imported by most molecule editors for conversion back into drawings or models of the molecules. Either the standard SMILES or the Canonical SMILES are given
InChI Identifier:	IUPAC I nternational C hemical I dentifier. This is a textual

	identification for chemical substances, that provides a standard, readable way of encoding molecular information and to facilitate the search for such information in databases and on the web.
Structure diagram available	Either Yes or No. If Yes is stated this will be a link to a separate window displaying the structure diagram.
Molecular mass:	The relative molecular mass (molecular weight) of a chemical is the mass of a molecule of the chemical relative to the mass of a carbon atom taken as exactly 12.
Chemical name:	Name of the chemical according to the nomenclature rules of IUPAC or CAS. Where this is not available or does not apply a generic name is given.
Other status information:	<p>This will display information relating to the status of the pesticide with respect to other legislation, international conventions and information regarding phase out. Including:</p> <p>PIC Annex 1 chemicals: The Rotterdam Convention on Prior Informed Consent (PIC). The Convention covers pesticides and industrial chemicals that have been banned or severely restricted for health or environmental reasons by parties and which have been notified for inclusion in the PIC procedure by those parties. One notification from each of two specified regions triggers consideration of addition of a chemical to Annex III of the Convention. Severely hazardous pesticide formulations that present a hazard under conditions of use in developing may also be nominated for inclusion in Annex III.</p> <p>POP chemicals: The Stockholm Convention on Persistent Organic Pollutants (POP). POPs are exceedingly toxic chemicals that are extremely persistent in the environment, travel long distances on wind and water currents, and concentrate up the food chain to accumulate in our bodies. They also have serious health implications and can cause cancer, neurological and learning disabilities, and subtle changes to human reproductive and immune systems. The Stockholm Convention bans or severely restricts the most hazardous POPs, and establishes an international, science-based process for adding other POPs to the treaty. Those listed in this database also include pesticides that are new POP candidates proposed by other organisations including the WWF.</p> <p>VOC chemicals: Volatile organic compounds (VOCs) are organic chemical compounds that have high enough vapor pressures under normal conditions to significantly vapourise and enter the atmosphere. The term VOC may have special legal meanings in some countries.</p> <p>LRTAP Chemicals: The Convention on Long-range Transboundary Air Pollution (LRTAP): The aim of the Convention is that Parties seek to limit and, as far as possible, gradually reduce and prevent air pollution including long-range transboundary air pollution. Chemicals considered to be the most serious problem are assigned to Annex 1.</p> <p>PAN Dirty Dozen / PAN Bad Actor: The Pesticide Action Network (PAN) have identified these chemicals as being particularly harmful.</p> <p>OSPAR: OSPAR Convention for the Protection of the Marine Environment: pfa - priority substances for action, soc - substances of concern.</p> <p>WFD: Water Framework Directive 2000/60/EC: phs - priority hazardous substance, pps - possible priority substance, other - other substance of concern.</p>

	Groundwater contaminant: substance is known to have polluted groundwaters and is a substance of concern.
Resistance code:	This is the HRAC, IRAC or FRAC code that denotes their resistance classification and can be used in resistance management programmes.
Physical state:	Provides an indication of the physical state of the material – solid, liquid or gas and its general appearance. This normally applies to the active substance in its pure state unless stated otherwise.

(2) General product and branding information

Parameter	Explanation
Example manufacturers &/or suppliers	Examples of companies who have manufactured, supplied or used the substance in their products.
Example products:	Non-exhaustive list of examples of products that include the substance.
Associated substances:	Non-exhaustive list of other substances associated with the substance. These may include other pesticides, solvents, inerts and impurities or concern. Links cross-referencing to the associated substances are supplied.
UK LERAP status:	Classification of the substance (where applicable) regarding its UK LERAP (Local Environmental Risk Assessment for Pesticides) status and the need for 'No spray' zones adjacent to surface water bodies.
Formulation & application details:	This provides brief information on the main types of formulations and aspects relating to their application.

(3) General chemical properties related to environmental fate

NB: where the stored parameter differs significantly from the information stated below this will be described in the accompanying text field.

Parameter	Explanation
Solubility in water: mg/l	The mass of a given substance (the solute) that can dissolve in a given volume of water. Value reported is at 20°C. Note for some pesticides solubility may be pH sensitive.
Solubility in organic solvents: mg/l	The mass of a given substance (the solute) that can dissolve in a given volume of solvent. Value reported is at 20°C. Note for some pesticides solubility may be pH sensitive.
Melting point: °C, at 1 atmosphere pressure	The temperature at which the given substance changes its physical state from solid to liquid.
Boiling point: °C, at 1 atmosphere pressure	The temperature at which the vapour pressure of the substance in its liquid state equals the environmental pressure surrounding the liquid i.e. it boils
Degradation temperature: °C, at 1 atmosphere pressure	The temperature at which the substance is no longer stable and begins to break down.
Flash point: °C	The flash point of a flammable substance is the lowest temperature at which it can form an ignitable mixture in air.
Octanol-water partition coefficient as LogP:	LogP is the logarithm (base-10) of the partition coefficient between n-octanol and water. It is used in environmental fate studies and large values (+4 or higher) are regarded as an indicator that a substance

	will bio-accumulate. For some substances LogP will be very sensitive to pH.
Bulk density/Specific gravity:	Parameter given depends on the physical state (solid or liquid) of the pesticide. Solids - Bulk Density is the weight of the pesticide per unit volume. Liquids - Specific Gravity is the ratio of the density of the pesticide to the density of water.
Dissociation constant pKa:	Strengths of acids and bases can be indicated on a common scale at 25°C. Defined as the negative logarithm of the acidity constant Ka. The lower the pKa the stronger the acid. For example acetic acid has a pKa of 4.75 whilst sulphuric acid has a pKa of -3.0. pKa is used here as an indicator of the potential of a compound to form ions in water. Many pesticide active substances are either permanently ionic or will change ionic state somewhere in the range of the pH of environmental soils and water. Knowing the ionic state of a pesticide provides important information on its potential mobility and persistence in the environment.
Vapour pressure: mPa	The pressure at which a liquid is in equilibrium with its vapour at 25°C. It is a measure of the tendency of a material to vapourise. The higher the vapour pressure the greater the potential.
Henry's law constant Dimensional / Dimensionless:	A Gas Law states that the amount of gas absorbed by a given volume of liquid at a given temperature is directly proportional to the partial pressure of that gas in equilibrium with that liquid. As such it provides an indication of the preference of a chemical for air relative to water i.e. its volatility. Henry's Law Constant is usually quoted in Pa.m ³ /mol or in a dimensionless form at 20°C.
DT50 Soil	DT50 is the time required for the pesticide concentration under defined conditions to decline to 50% of the amount at application. In many cases pesticides show "half-life" behaviour, in which subsequent concentrations continue to decline by 50% in the same amount of time. In such cases several or more four half-lives (in which the concentration declines to 1/8 or 1/16, e.g.) are a measure of the persistence on the pesticide.
DT90 Soil	DT90 is the time required for the pesticide concentration under defined conditions to decline to 90% of the amount at application.
Soil degradation DT50 Typical/ Laboratory/ Field: (days)	DT50 in a field or laboratory soil sample. Three parameters are given. Typically data is derived from laboratory studies, but where the substance is persistent in soil under laboratory conditions, field studies may be carried out. 'Typical values' quoted are those given in the general literature and are often a mean of all studies field and laboratory. This is the value normally used in the regulatory modelling studies and is for aerobic conditions.
Soil degradation DT90 Typical/ Laboratory/ Field: (days)	DT90 in a field or laboratory soil sample. Three parameters are given. Typically data is derived from laboratory studies, but where the substance is persistent in soil under laboratory conditions, field studies may be carried out. 'Typical values' quoted are those given in the general literature and are often a mean of all studies field and laboratory. Data is for aerobic conditions.
Aqueous photolysis DT50 (days) at 20°C pH 7:	Photochemical processes may be important in determining the fate of organic pollutants in aqueous environments. This is the rate of chemical decomposition in the aquatic environment induced by light or other radiant energy expressed as a DT50. Other information regarding, for example, pH sensitivity is also given in accompanying notes.

Aqueous hydrolysis DT50 (days) at 20°C and pH 7:	This is the rate of chemical decomposition induced by water at pH 7 expressed as a DT50. Other information regarding, for example, pH sensitivity is also given in accompanying notes.		
Water sediment study data: Water-sediment & Water phase only DT50s:	This is the rate of chemical decomposition in water-sediment systems expressed as a DT50. Data is given for the system as a whole and for the water phase only.		
GUS leaching potential index: NOTE: THIS IS AN INDICATOR & NOT A RISK ASSESSMENT	<p>The GUS index (Groundwater Ubiquity Score) is a very simple indicator of a chemical potential for leaching into groundwater. It is based on the environmental fate properties of the chemical and takes no account of environmental conditions. It is not a substitute for modelling and risk assessment studies.</p> <p>Calculated from the soil degradation rate (DT50) and the Organic-carbon sorption constant (koc) where: $GUS = \log(DT50) \times (4 - \log(koc))$</p> <p>If $GUS > 2.8$ = pesticide likely to leach If $GUS < 1.8$ = pesticide unlikely to leach If $GUS 1.8 - 2.8$ = leaching potential is marginal</p> <p>(Reference: Gustafson, D.I. (1989) Groundwater Ubiquity Score: A Simple Method for Assessing Pesticide Leachability Environmental Toxicology and Chemistry, 8, pp339-357).</p>		
SCI-Grow groundwater index: NOTE: THIS IS AN INDICATOR & NOT A RISK ASSESSMENT	<p>This is an indicator, used by the USEPA, to crudely estimate pesticide concentrations in vulnerable groundwater. It is based on environmental fate properties of the pesticide, the application rate and existing data from small-scale monitoring studies. It is not a substitute for modelling and risk assessment studies.</p> <p>(Ref: www.epa.gov/oppefed1/models/water/scigrow_description.htm)</p> <p>Calculated from the soil degradation rate (DT50) and the Organic-carbon sorption constant (koc) where:</p> <p>Step 1:</p>		
	D = $\log(Koc + 5.0)$		
	if DT50 < 6.0 then	C = $\log(DT50/6.0)$	R = C * D
	if DT50 6.0 - 1500 then	C = $\log(DT50 - 5.0)$	R = C * (4-D)
	if DT50 > 1500 then	C = $\log(1500) = 3.176$	R = C * (4.-D)
	Step 2:		
	If Koc = 9995 then	F = $-2.241 + 0.610 * R$	Normalised concentration = 10^F
	If Koc > 9995 then	Normalised concentration = 0.006	
	<p>Step 3:</p> <p>Total pesticide use = application rate (kg/ha) * number of applications</p> <p>Estimated concentration in groundwater = Normalised concentration * total pesticide use</p> <p>(Reference: www.oecd.org/dataoecd/6/17/2752913.pdf)</p>		
Potential for particle bound transport index: NOTE: THIS IS AN INDICATOR & NOT A RISK ASSESSMENT	This is a simple indicator of a pesticides run-off potential. It classifies pesticides into three broad risk categories based on environmental fate properties. It takes no account of environmental or landscape conditions. It is not a substitute for modelling and risk assessment studies.		
Koc (Kfoc) - organic-	Chemicals vary in how well they are adsorbed to soil particles. Koc /		

carbon sorption constant: (mL/g)	K _{foc} measures the affinity for pesticides to sorb to organic carbon. The higher the value, the stronger the tendency to attach to and move with soil. K _{oc} / K _{foc} values greater than 1000 indicate strong adsorption to soil. Chemicals with lower K _{oc} / K _{foc} values (less than 500) tend to move more with water than be adsorbed to sediment. Some pesticides are strongly bound to other soil components such as clay surfaces. For some pesticides K _{oc} / K _{foc} will be very pH sensitive.
Freundlich coeffs (K _f and ^{1/n}):	A 'soil absorption coefficient' study is to predict the mobility of a pesticide in the environment. If carried out over a range of pesticide concentrations the result can be fitted to an equation called the 'Freundlich isotherm', that is used to describing adsorption. This equation is $Q=kC^{1/n}$, where Q is the amount of the sorbed pesticide in equilibrium, C is the concentration of the pesticide in solution, K and n are empirical constants, K _f is called the sorption coefficient. An n value of 1 denotes a linear sorption process.
UV-Visible spectrum (L mol ⁻¹ cm)	The specific wavelengths of light absorbed by a molecule depend on the types of functional groups and this information can help with qualitative identification of these functional groups.

(4) Metabolite/Parent information

Key soil metabolites:

Parameter	Explanation
Metabolites:	A pesticide degradation product - a chemical product formed upon the breakdown of a pesticide or other chemical. Any significant or important metabolites are shown (predominately those formed in the soil) together with links to their profile pages. This list is not exhaustive. Links cross-referencing to the metabolites are supplied.
Parent:	The parent substance that has degraded to produce the metabolite.
Formation media:	The host process that creates the metabolite e.g. soil, water, animal, plant, groundwater.
Estimated Maximum Occurrence:	The maximum amount of the named metabolite identified (via soil degradation studies) in the formation media expressed as a fraction of the pesticide applied.
91/414 Relevancy	<p>Within Directive 91/414 a metabolite is considered to be 'relevant':</p> <ul style="list-style-type: none"> - if there is reason to believe that it has comparable intrinsic properties to the parent chemical in terms of its biological target activity, or - if it - has toxicological properties that are considered severe, or - poses a higher or comparable risk to organisms than the parent substance. <p>Therefore 'relevant' metabolites may need to so be taken into account within any risk assessment process. Please note that some 91/414 dossiers pre-date the introduction of 'relevant metabolites' and, instead, declared metabolites as 'Major' if their estimated maximum occurrence fraction >10%.</p>

Other metabolites:

Parameter	Explanation
Chemical name	The chemical name using the CAS nomenclature system
Alias	Other common names by which the metabolite may be known.
Ref code	The manufacturers reference Identifier
Formation medium	The media in which the substance can be found
Estimated Maximum Occurrence:	The maximum amount of the named metabolite identified (via soil degradation studies) in the formation media expressed as a fraction of the pesticide applied.
Other notes	Other data related to the metabolite.

(5) Ecotoxicology

Parameter	Explanation
Bio-concentration factor:	The concentration of the pesticide in tissue per concentration of chemical in water. This describes the accumulation of pollutants through chemical partitioning from the aqueous phase into an organic phase, such as the gill of a fish.
Bioaccumulation potential:	The potential for an organism to absorb the pesticide at a rate greater than that at which the substance is lost from the body.
Fauna and flora ecotoxicological end point data:	Data used to measure the adverse effects on living organisms that chemicals can have when released into the natural environment. See Endpoint Glossary and Key Species data below.
Mesocosm study data:	Mesocosm studies are a useful tool for higher-tier aquatic risk assessment. They are widely used in the pesticide regulations process to evaluate the effects of pesticides on aquatic communities at the ecosystem level.

(6) Human health

Parameter	Explanation
Toxicity endpoints for mammals:	Oral, dermal and inhalation values provided.
WHO classification:	Toxicity hazard class given by the World Health Organisation (WHO). Class Ia: extremely hazardous Class Ib: highly hazardous Class II: moderately hazardous Class III: slightly hazardous O: Obsolete substance NL: Not listed The system is based on the LD50 endpoint for rats. An ingested solid with a LD50 5mg or less/kg bodyweight is Class Ia, at 5-50 mg/kg Class Ib, at 50-500 mg/kg Class II, and at more than 500 mg/kg Class III. Values may differ for liquid oral agents and dermal agents.
US EPA acute toxicity class (formulation)	Toxicity hazard class of the formulation as given by the EPA. Class 1: Highly toxic Class II: Moderately toxic Class III: Slightly toxic Class IV: Not acutely toxic

US EPA classification for products:	This is the hazard class assigned to pesticide products containing the active substance. Generally, Class I substances will kill an adult person at a dose of less than 5g, Class II at 5-30g, and Class III >30g.
EC risk classification / safety classification:	EC Directive 67/548/EEC provides a harmonised basis for the classification (risk) and handling (safety) of dangerous substances. These listed here are those for the active substance. More details of what the specific codes means can be found on the PPDB website, in the downloads section.
International Maritime Dangerous Goods (IMDG) Code	The IMDG Code is an international code related to the transport of dangerous goods by sea covering packing, container traffic and stowage. Dangerous goods are categorised into different classes, further subdivided to define and describe characteristics and properties of the substances which would fall within each class.
UN Packaging Mark	The packaging used to transport dangerous substances normally has to meet certain standards defined by the United Nations (UN). The UN marking system indicates several characteristics of the packaging, the test levels the packaging has successfully passed and provides an indication of any hazardous properties the material packed in the container may exhibit.
Acceptable Daily Intake – ADI mg kg ⁻¹ bw day ⁻¹ :	The amount of chemical that can be consumed each day without causing harm - as far as evidence suggests.
Acute Reference Dose – ARfd mg kg ⁻¹ bw day ⁻¹ :	The amount of chemical that can be ingested over a short period of time, usually during one meal or one day, without appreciable health risk to the consumer - as far as evidence suggests.
Acceptable Operator Exposure Level mg kg ⁻¹ bw day ⁻¹ : - AOEL:	This is a health-based limit that is established on the basis of the full toxicological assessment required for pesticide registration. The risk for pesticide operators can be quantified by comparing this value with exposure level during pesticide application.
Dermal penetration studies %	Mean value of known studies reported. Default value in the absence of further evidence is normally taken as 10%.
Dangerous substances Directive:	This Directive concerns pollution caused by certain dangerous substances discharged into the aquatic environment and aims to regulate potential aquatic pollution. The Directive covered discharges to inland surface waters, territorial waters, inland coastal waters and groundwater. The protection of groundwater is now regulated under a separate Council Directive. Directive 76/464 introduced the concept of list I and list II chemicals. Where the pesticide concerned appears on these lists it is given here.
Exposure limits:	Exposure of employees, operators and other workers to hazardous substances should be prevented and in some cases national or EC regulations impose limits. Where identified these are quoted and may include Maximum Exposure Limits (MEL), Occupational Exposure Standards (OES) and Short Term Exposure Limits (STEL). List is not exhaustive but given for guidance only.
Exposure routes public and occupational:	Where identified risks to the public (e.g. bystanders and consumers) and operators/workers are identified. This list is not exhaustive but given for guidance only.

EU MRLs mg kg ⁻¹	MRLs are defined as the maximum concentration of pesticide residue likely to occur in or on food, drink and feeding stuffs after the use of pesticides according to Good Agricultural Practice (GAP). GAP is defined as the pesticide being applied in accordance with the product label recommendations and in keeping with local environmental and other conditions. The values listed here are for guidance only and are often those <u>proposed</u> within the EFSA DAR and Conclusion documents. More precise information can be found in the EU Database .
Drinking water MAC:	The Maximum Acceptable Concentration of the chemical in drinking water.
General health issues:	Summary of main issues relating to human health. This list is not exhaustive.
Specific health issues:	Summary of the main human health concerns across a number of issues. Note this is somewhat subjective as literature is not universally uniform in the way issues are addressed. We have used a 'weight-of-the-evidence' approach erring on the side of caution.
Handling issues:	General description of any hazards that should be addressed when handling the substances.

(7) Translations

Language translations for the common name by which the substance is known is provided (where known) in English, French, German, Danish, Italian, Spanish, Greek, Slovenian, Polish, Swedish, Hungarian and Dutch.

The Primary Data Sources Used to Populate the PPDB

Note: This list is non-exclusive and just provides an example of the sources utilised

A	EU Regulatory & Evaluation Data as published by EC and EFSA (DAR & Conclusion dossiers) / EU Annex III PIC DGD / EU MRL Database (See http://ec.europa.eu/sanco_pesticides/public/index.cfm)
AA	IOBC Database on classification of side effects to beneficial organisms, 2005
AB	SELECTV Database (See http://ipmnet.org/phosure/database/selctv/selctv.htm)
AC	EC Joint Research Centre ESIS European Chemical Substance Information Systems including EINECS (See http://ecb.jrc.ec.europa.eu/esis/)
AE	Joint Assessment of Commodity Chemicals ECETOC (See http://www.ecetoc.org)
AF	European Food Safety Agency (EFSA)
B	UK PSD ACP Evaluation Documents / and other DEFRA (UK) documents (See http://www.pesticides.gov.uk/publications.asp?id=202)
C	AGRITOX (See http://www.dive.afssa.fr/agritox/index.php)
CA	Medical and toxicological databases and information systems e.g. TOXNET (See http://toxnet.nlm.nih.gov/cgi-bin/sis/htmlgen?HSDB)
D	Agricultural Research Information System (ARIS) Database
DW	Don Wauchope personal database for Pka data: Wauchope, R. D. and Edwards, J. Dissociation constants for pesticide active ingredients: a database and comparison with predicted values. MS in preparation
E	Manufacturers Safety Data Sheets

F	U.S. EPA ECOTOX Database (see http://cfpub.epa.gov/ecotox/) / U.S. EPA Pesticide Fate Database (See http://cfpub.epa.gov/pfate/home.cfm) / Miscellaneous WHO documents.
FAO	Miscellaneous FAO publications
G	Extension Toxicology network Database EXTOWNET (See http://extownet.orst.edu/ghindex.html)
H	The US ARS Pesticide Properties Database (See http://www.ars.usda.gov/Services/docs.htm?docid=14199)
J	Pesticide Action Network Database (See http://www.pesticideinfo.org/)
K	Research Datasets (e.g. Pandora, Demetra (see http://www.demetra-tox.net/))
L	Pesticide manuals and hard copy reference books / other sources
M	GLEAMS Model database (Groundwater Loading Effects of Agricultural Management Systems). (See http://www.cpes.peachnet.edu/sewrl/Gleams/gleams_y2k_update.htm)
N	Various Trusts, NGOs & Charities Data
P	Other Governments and Regulators
O	Miscellaneous Data From On-line Sources
R	Peer Reviewed Scientific Publications
S	Expert Judgement
T	UN EPFA Database
US	US Dept of Agriculture National Resources Conservation Service - various datasheets, databases and online sources
V	ChemID Online Databases (See http://chem.sis.nlm.nih.gov/chemidplus/) / IPCS INCHEM (See http://www.inchem.org/)
W	French database provided by ARVALIS-Institut du Végétal
X	WINPST Database (See http://www.ipm.ucdavis.edu/TOX/winpstdoc.html)
Y	Germany's Federal Environment Agency (UBA) (See http://www.umweltbundesamt.de/index-e.htm)
Z	Kingtai Chemicals Website (See http://www.kingtaichem.com/)

Data Interpretation

(1) Environmental fate

Parameter	Source	Thresholds
Solubility in water (mg l ⁻¹):	Arbitrary system in common use	<= 50 = Low 50 - 500 = Moderate > 500 = High
Octanol-water Partition Coefft (Log P):	Used by the US EPA. Widely used rule of thumb.	< 2.7 = Low bioaccumulation 2.7 - 3 = Moderate > 3.0 = High
Soil degradation (days):	See note 1.	< 30 = Non-persistent 30 - 100 = Moderately persistent 100 - 365 = Persistent > 365 = Very persistent
Aqueous photolysis DT50 (days) at pH 7:	See note 5.	< 1 = Fast 1 - 14 = Moderately fast 14 - 30 = Slow > 30 = Stable
Aqueous hydrolysis DT50 (days) at 20°C and pH 7:	See note 5.	< 30 = Non-persistent 30 - 100 = Moderately persistent 100 - 365 = Persistent > 365 = Very persistent
Water-sediment degradation (days):	Same thresholds as soil degradation utilised. See note 1.	< 30 = Fast 30 - 100 = Moderately fast 100 - 365 = Slow > 365 = Stable
Water phase only degradation (days):	See note 5.	< 1 = Fast 1 - 14 = Moderately fast 14 - 30 = Slow > 30 = Stable
GUS Index:	Gustafson, DI (1989) Groundwater Ubiquity Score: a simple method for assessing pesticide leachability. Environ. Toxicol. Chem 8, 339-357.	> 2.8 = High leachability 2.8 - 1.8 = Transition state < 1.8 = Low leachability
Koc/ Kfoc (ml g ⁻¹):	PSD Pesticide Data Requirement Handbook (2005). SSLRC Mobility Classification System. Also See note 5.	< 15 = Very mobile 15 - 75 = Mobile 75 - 500 = Moderately mobile 500 - 4000 = Slightly mobile > 4000 = Non-mobile
Vapour pressure at 25°C (mPa):	Kerle EA, Jenkins JJ & Vogue PA (1996) Understanding pesticide persistence and mobility for groundwater and surface water protection. Oregon State University. EM 8561.	< 1 x 10 ⁻⁶ = Non-volatile 1 X 10 ⁻⁴ - 1 x 10 ⁻⁶ = Intermediate state > 1 X 10 ⁻⁴ = Volatile
Henry's Law Constant at 20°C (Dimensionless - K _H):	See note 2.	> 2.5 x 10 ⁻⁵ = Volatile 2.5 x 10 ⁻⁷ - 2.5 x 10 ⁻⁵ =

		Moderate volatility < 2.5×10^{-7} = Non volatile
Henry's Law Constant ($\text{Pa m}^3 \text{ mol}^{-1}$):	Rule of thumb in wide, general use.	> 100 = Volatile 0.1 - 100 = Moderately volatile < 0.1 = Non-volatile
Particle Bound Transport Indicator:	Goss & Wauchope (1990). See note 3.	
SCIGROW Indicator:	See note 10.	
Freundlich equation:	See note 11.	

(2) Ecotoxicology

Parameter	Source	Thresholds
Bio-concentration factor:	General rule of thumb and that used by the US EPA.	< 100 = Low potential 5000 - 100 = Threshold for concern > 5000 - High potential
Bioaccumulation potential:	See note 7.	See note 7.
Mammals - Acute oral LD50 (mg kg^{-1}):	See note 5.	> 2000 = Low 100-2000 = Moderate < 100 = High
Mammals - Short term dietary NOEL (mg kg^{-1}):	See note 5.	> 2000 = Low 100-2000 = Moderate < 100 = High
Birds - Acute LD50 (mg kg^{-1}):	Consistent with US EPA Guidelines. See note 5.	> 2000 = Low 100 - 2000 = Moderate < 100 = High
Fish - Acute 96 hour LC50 (mg l^{-1}):	See note 5.	> 100 = Low 0.1 - 100 = Moderate < 0.1 = High
Fish - Chronic 21 day NOEC (mg l^{-1}):	See note 5.	> 10 = Low 0.01 - 10 = Moderate < 0.01 = High
Aquatic invertebrates - Acute 48 hour EC50 (mg l^{-1}):	See note 5.	> 100 = Low 0.1 - 100 = Moderate < 0.1 = High
Aquatic invertebrates - Chronic 21 day NOEC (mg l^{-1}):	See note 5.	> 10 = Low 0.01 - 10 = Moderate < 0.01 = High
Aquatic crustaceans - Acute 96 hour LC50 (mg l^{-1}):	See note 5.	> 100 = Low 0.1 - 100 = Moderate < 0.1 = High
Sediment dwelling organisms - Acute 96 hour LC50 (mg l^{-1}):	See note 5.	> 100 = Low 0.1 - 100 = Moderate < 0.1 = High
Sediment dwelling organisms - Chronic 28 day NOEC, static, water (mg l^{-1}):	See note 5.	> 10 = Low 0.01 - 10 = Moderate < 0.01 = High
Sediment dwelling organisms - Chronic 28 day NOEC, sediment (mg kg^{-1}):	See note 5..	> 100 = Low 0.1 - 100 = Moderate < 0.1 = High
Aquatic plants - Acute 7 day	See note 6.	> 10 = Low

EC50, biomass (mg l ⁻¹):		0.01 - 10 = Moderate < 0.01 = High	
Algae - Acute 72 hour EC50, growth (mg l ⁻¹):	See note 6.	> 10 = Low 0.01 - 10 = Moderate < 0.01 = High	
Algae - Chronic 96 hour NOEC, growth (mg l ⁻¹):	See note 6.	> 1 = Low 0.001 - 1 = Moderate < 0.001 = High	
Honeybees - Acute 48 hour LD50 (µg bee ⁻¹):	See note 5.	> 100 = Low 1 - 100 = Moderate < 1 = High	
Earthworms - Acute 14 day LC50 (mg kg ⁻¹):	See note 12.	> 1000 = Low 10 - 1000 = Moderate < 10 = High	
Earthworms - Chronic 14 day NOEC, reproduction (mg kg ⁻¹):	See note 12.	> 100 = Low 0.1 - 100 = Moderate < 0.1 = High	
Other soil macro-organisms:	LR50 / EC50 / NOEC / % Effect:	See note 13.	No interpretation
Other arthropods:	LR50 data:	See note 8.	< 50 = Harmful at 1kg ha ⁻¹ 50-500 = Moderately harmful at 1kg ha ⁻¹ > 500 = Harmless at 1kg ha ⁻¹
	% Effect data:	See note 9.	< 30 = Harmless 30 - 79 = Moderately harmful > 79 = Harmful
Soil micro-organisms:	Nitrogen / carbon mineralisation.		<= 25% change is considered insignificant.

(3) Human health

Parameter	Source	Thresholds
Who Toxicity class:	WHO Guidelines 2004. Based on rat LD50 & physical state of the pesticide. See note 4.	See note 2.
EPA acute toxicity class (formulation):	Standard text book thresholds. These apply to the formulated product.	I - Highly toxic II - Moderately toxic III - Slightly toxic IV - Not acutely toxic In some instances there is slight variation in product classification. In these instances multiple listings are given, unless there is no consensus across brands in which case the text reads 'No consensus across products'.
EC Risk classification:	Risk Information contained with in Annex II and III of Directive 67/548/EEC on Classification and Labelling of Dangerous Substances.	

EC Safety classification:	Safety Information contained with in Annex IV of Directive 67/548/EEC on Classification and Labelling of Dangerous Substances.
ADI (mg kg ⁻¹ bw):	The acceptable daily intake is the amount of a substance that can be ingested every day of an individual's entire lifetime, in the practical certainty, on the basis of all known facts, that no harm will result. SF – refers to the safety factor applied.
AOEL (mg kg ⁻¹):	The acceptable operator exposure level is the maximum amount of active substance to which the operator may be exposed without any adverse health effects. SF – refers to the safety factor applied.
Dangerous substances directive:	This Directive requires Member States to introduce measures to eliminate (List I) or to reduce (List II) pollution of the aquatic environment from certain listed substances identified in its Annexes.
Exposure limits:	An exposure limit is the concentration of a chemical in the workplace air to which most people can be exposed without experiencing harmful effects.
MRL's - maximum residue limits:	These limits can change and the data given here is usually that proposed by EFSA. Data may not be complete. See EU database for further information.
Drinking water MAC:	Maximum Admissible Concentration of the pesticide in drinking water. The MAC for a pesticide is derived from its ADI. The EU Drinking Water Directive imposes a maximum admissible concentration (EU MAC) for any individual pesticide compound of 0.1 mg l ⁻¹ .

Notes

1. Consistent with EU Guidance. (9188/VI/97 rev. 8.) and
 - I. Kerle EA, Jenkins JJ & Vogue PA (1996), Understanding pesticide persistence and mobility for groundwater and surface water protection. Oregon State University. EM 8561.
 - II. Rao PSC & Hornsby AG (2004) Behaviour of pesticides in Soils and water. University of Florida. See <http://edis.ifas.ufl.edu/SS111>.
 - III. See also Note 3 below.
2. Several relevant references which include:
 - I. Van der Werf , HMG (1996) Assessing the impact of pesticides on the environment. Agriculture, Ecosystems & Environment, 60, 81-96.
 - II. Jury WA, Spencer WF, & Farmer WJ (1984) Behaviour assessment model for trace organics in soil. III Application of screening model. J. Environ Qual. 13, 573-579.
 - III. Kerle EA, Jenkins JJ & Vogue PA (1996) Understanding pesticide persistence and mobility for groundwater and surface water protection. Oregon State University. EM 8561.
3. Table below has been extracted from:
 - I. Goss, D & Wauchope RD (1990) The SCR/ARS/CES Pesticide Properties Database. II using it with Soils data in a screening Procedure. In D.L. Weigmann Ed., Pesticides in the next decade: the challenge ahead, Virginia Resources Research Centre, Blacksburg, VA, USA pp471-493.

Potential for Particle-bound transport	Criteria
High	DT50 \geq 40 days & Koc \geq 1000 DT50 \geq 40 days, Koc \geq 500 & solubility = 0.5 mg/l
Low	DT50 \leq 1 day DT50 \leq 2 days & koc \leq 500 DT50 \leq 4 days, Koc \leq 900 & solubility \geq 0.5 mg/l DT50 \leq 40 days, Koc \leq 500 & solubility \geq 0.5 mg/l DT50 \leq 40 days, Koc \leq 900 & solubility \geq 2 mg/l
Medium	All other

4. Classification given below has been extracted from the WHO Guidelines document: The WHO recommended classification of pesticides by hazard & guidelines to classification. (2004). See <http://www.who.int/publications/en/>

Class Ia: extremely hazardous
Class Ib: highly hazardous
Class II: moderately hazardous
Class III: slightly hazardous
O: Obsolete
NL: Not listed

5. Thresholds used have been selected to be consistent with industry guidelines, were developed, and are consistent with regulatory thresholds used in both the UK and EU.
6. The EU (Uniform Principles) (Annex VI of Directive 91/414/EEC) guidelines have been adopted have set toxicity:exposure (TER) ratios for algae and aquatic plants at $\frac{1}{10}$ th of those for fish and daphnids. The same ratio has been applied here.
7. Deleted
8. In EU pesticide regulatory risk assessments 'hazard quotients' are used to determine the need for additional studies to assess risk to beneficial arthropods. Hazard quotients (HQ) are determined by dividing the application rate of the active substance by the median lethal rate (LR50). HQ values less than 2.0 are considered to be low risk to beneficial arthropods and additional (higher tier) data is not required. Values greater than 2.0 trigger additional data requirements. The risk phrases quoted within this database are based on a standard application rate of 1 kg ha⁻¹ and should be taken as **a comparative guideline only**. For a more accurate assessment divide the actual application rate of the active substance (in grams) by the LR50 value quoted. Any value less than 2.0 can be considered a low risk. Values greater than 2.0 may require care and mitigation. Please note the thresholds for moderate and high risk are **arbitrary values**.
9. Where two values are given (e.g. mortality / parasitism), base assessment on the highest absolute value (i.e. can be a positive or negative effect).
10. SCI-GROW is a screening model used by the US EPA to estimate pesticide concentrations in vulnerable groundwater. The model provides an exposure value that can be used to determine the potential risk to the environment and to human health from drinking water contaminated with the pesticide. The SCI-GROW estimate is based on environmental fate properties of the pesticide (aerobic soil degradation half-life and linear adsorption coefficient normalised for soil organic carbon content), the maximum application rate, and

existing data from small-scale prospective ground-water monitoring studies at sites with sandy soils and shallow ground water.

SCI-GROW estimates represent worse case estimates. For this reason, it is not appropriate to use SCI-GROW concentrations for national or regional exposure estimates. Nor is this indicator an alternative to a scientific risk assessment. Values given are based on a standard 1 kg ha⁻¹ or 1 L ha⁻¹ application rate and should be adjusted to the actual application rate used

For more information see:

http://www.epa.gov/oppefed1/models/water/scigrow_description.htm.

11. The distribution of a pesticide between the solution and adsorbed phases can often be described by the "Freundlich equation", an equation that is used to describe a wide variety of adsorption data from every area of science. The equilibrium concentration and adsorbed pesticide amounts are determined experimentally. The Log₁₀ of the quantity of adsorbed pesticide is plotted against the equilibrium concentrations. Often the relationship obtained is approximately linear and can be described by the Freundlich equation: $Q = KC^{1/n}$, where Q is the adsorbed amount of pesticide (μg kg⁻¹), C is the equilibrium concentration (μg l⁻¹), and kf and n are the experimental parameters unique to the isotherm. The parameter n is greater than 1, the larger it is the more non-linear the equation becomes.
12. The availability of the pesticide in the soil can depend on the amount of soil organic carbon (SOC). The toxicity endpoint value may therefore be corrected for the difference in SOC of the test soil and the reference soil. This means that the toxicity endpoint value is divided by the percentage organic matter in the standard test soil, and multiplied by the percentage organic matter in the reference soil. Uncorrected values are quoted herein unless otherwise stated e.g. '(corr)'.
13. Data is very limited and is presented in the literature in a variety of formats. Therefore neither a standard format nor interpretation can be provided.

Key Species and Endpoint Glossary

Parameter	Explanation
EC50:	The concentration of a chemical that can be expected to cause a defined non-lethal effect in 50 per cent of the tested population. In some cases other percentages may be displayed e.g. EC10 or EC15.
LD50:	Used in toxicology this is the median lethal dose (LD ₅₀ , abbreviation for 'Lethal Dose, 50%'), of a toxic substance and is the dose required to kill half the tested population. LD ₅₀ figures are frequently used as a general indicator of a substance's acute toxicity.
NOEL/NOEC:	Greatest concentration or level of a substance, found by observation or experiment, which causes no detectable effect.
NOEAEC:	'No Observed Ecologically Adverse Effect Concentration' - the highest concentration that causes no observed adverse effect on fauna or flora.

Species	Explanation
Given below are the first choice species for ecotoxicological endpoints. However, if these are not available data for other species may be given instead. This will be indicated in the accompanying notes. Where data for several species are available, data for the most sensitive is given. (See also 'PPDB Conditions for Use' document)	
Mammals	Rat, Mice, Dog (Mainly used for the human health studies)
Birds	Mallard duck (<i>Anas platyrhynchos</i>) Bobwhite quail (<i>Colinus virginianus</i>)
Fish	Rainbow trout (<i>Oncorhynchus mykiss</i>) Bluegill sunfish (<i>Lepomis macrochirus</i>) Zebra fish (<i>Brachydanio rerio</i>)
Aquatic invertebrates	Daphnids (<i>Daphnia magna</i> , <i>Daphnia pulex</i>)
Aquatic crustaceans	Mysid shrimps (<i>Americamysis bahia</i>)
Sediment dwellers	Chironomid midges (<i>Chironomus riparius</i>)
Higher aquatic plants	Duckweed (<i>Lemna gibba</i> , <i>Lemna minor</i>)
Algae	Green algae (<i>Pseudokirchneriella subcapitata</i> , <i>Scenedesmus subspicatus</i>)
Honeybees	European honeybee (<i>Apis mellifera</i>)
Soil macro-organisms	Common brandling worm (<i>Eisenia foetida</i>)
Other soil dwelling arthropods	Springtails (<i>Folsomia candida</i>)
Other arthropods	Aphid parasitoid (<i>Aphidius rhopalosiphi</i>) Predatory mite (<i>Typhlodromus pyri</i>)

Species	Explanation
For ecotoxicological data the 'worst case' data has been selected unless it appears wildly out of character with the majority of studies published. We have chosen specific species and endpoints where ever possible to ensure a harmonised and balanced data set. In some cases endpoints other than these may be used, where this is the case the accompanying text will provide additional information.	
Mammals	acute oral LD ₅₀ and short term 90-day NOEL as mg kg ⁻¹ bw
Birds	acute oral LD ₅₀ as mg kg ⁻¹ bw

Fish	acute 96-hr LC ₅₀ and 21day NOEC as mg l ⁻¹
Aquatic invertebrates	acute 48-hr ErC ₅₀ and 21day NOEC as mg l ⁻¹
Aquatic crustaceans	acute 96hr LC ₅₀ as mg l ⁻¹
Sediment dwellers	96-hr LC ₅₀ and chronic 28-day NOEC static water only and sediment as mg l ⁻¹
Higher aquatic plants	14-day EC ₅₀ as mg l ⁻¹
Algae	acute 72-96-hr EC ₅₀ and 72-96-hr NOEC biomass / growth as mg l ⁻¹
Honeybees	Lowest or contact and oral LD ₅₀ as ug/bee
Soil macro-organisms	14-day LC ₅₀ and chronic reproduction NOEC as mg kg soil ⁻¹ dw
Other soil dwelling arthropods	various endpoints depending on availability as mg l ⁻¹
Other arthropods	various endpoints depending on availability
Carbon & nitrogen mineralisation	% effects.

Quality Control and Management

Data management activities involve trawling for new or previously unidentified data. Evaluating its fitness-for-purpose, undertaking quality control activities and adding it to the database. The database and website are updated several times each week.

Data is sourced globally from published scientific literature and databases, manuals, registration databases and dossiers, company technical datasheets and research projects.

Prior to entry into the PPDB data is subject to quality control. This involves peer review, cross checking against other databases and data sources and, where doubt exists and the original reference is known, the original source is revisited.

Fitness for purpose is evaluated based on experimental conditions being appropriate for Europe, scientific protocols utilised, quality control findings and how recent the data is.

Data is then weighted 1 (low) to 5 (high) according to the confidence we have in that data. A low score does not necessarily indicate incorrect data but indicates we have not been able to obtain verification. Generally, as a guide the weighting scores are assigned according to the following:

1	Estimated data with little or no verification
2	Unverified data of unknown source
3	Unverified data of known source
4	Verified data
5	Verified data used for regulatory purposes.

However, confidence weightings may be adjusted if reason to doubt verified data exists or there is reason for greater confidence in an unverified value.

Once data has been accepted for inclusion into the database, data is extracted and transformed with respect to units. References are recorded and hardcopies filed.

Contact details:

PPDB Management Team
Agriculture and Environment Research Unit, Science & Technology Research Institute
University of Hertfordshire
College Lane
Hatfield, Herts., AL10 9AB, UK

Telephone: +44 (0)1707 284548
Email: aeru@herts.ac.uk